

$\lambda\mu$ -calculus and duality: call-by-name and call-by-value

Jérôme Rocheteau

ESTAS – INRETS
20 rue Élisée Reclus – BP 317
F-59666 Villeneuve d'Ascq Cedex
jerome.rocheteau@inrets.fr

Preuves, Programmes et Systèmes
CNRS – Université de Paris VII
UMR 7126 – Case 7014
175 rue du Chevaleret – 75013 Paris – France

Abstract. Under the extension of Curry-Howard's correspondence to classical logic, Gentzen's NK and LK systems can be seen as syntax-directed systems of simple types respectively for Parigot's $\lambda\mu$ -calculus and Curien-Herbelin's $\bar{\lambda}\mu\tilde{\mu}$ -calculus. We aim at showing their computational equivalence. We define translations between these calculi. We prove simulation theorems for an undirected evaluation as well as for call-by-name and call-by-value evaluations.

1 Introduction

Key systems for classical logic in proof theory are Gentzen's NK and LK. The logical equivalence between the latter was proved in [Gentzen, 1934]. We deal with the extension of Curry-Howard's correspondence between proofs and programs through the systems of simple types for the $\lambda\mu$ and $\bar{\lambda}\mu\tilde{\mu}$ -calculi. This extension concerns some other calculi. It is initially Felleisen's λc -calculus. Its type system is the intuitionistic natural deduction with the double negation axiom. Griffin proposed this axiom as the type for the c -operator in [Griffin, 1990]. However, we focus on calculi that correspond closer to Gentzen's systems. The $\lambda\mu$ -calculus was defined for NK in [Parigot, 1992]. The $\bar{\lambda}\mu\tilde{\mu}$ -calculus was designed for LK in [Curien and Herbelin, 2000]. In the general case, these two calculi are not deterministic. There exists critical pairs. The $\bar{\lambda}\mu\tilde{\mu}$ -calculus admits two deterministic projections depending on choosing one of the two possible symmetric orientations of a critical pair. They correspond to the call-by-name/call-by-value duality.

We aim at proving the computational equivalence between $\lambda\mu$ and $\bar{\lambda}\mu\tilde{\mu}$ -calculi. A major step was reached with the proof of the simulation of the $\lambda\mu$ -calculus by the $\bar{\lambda}\mu\tilde{\mu}$ -calculus in [Curien and Herbelin, 2000]. It holds both for call-by-name and call-by-value evaluations. We present the call-by-name/call-by-value projections of the $\lambda\mu$ -calculus in the same way as for the $\bar{\lambda}\mu\tilde{\mu}$ in [Curien and Herbelin, 2000]. It consists of choosing one of the two possible orientations of a critical pair. We prove that the $\lambda\mu$ -calculus simulates backwards the $\bar{\lambda}\mu\tilde{\mu}$ -calculus in such a way that we obtain easily the same result for the call-by-name, for the call-by-value and for the simple type case. The $\bar{\lambda}\mu\tilde{\mu}$ -calculus is composed of three syntactic categories: terms, contexts (or environments) and commands. The $\lambda\mu$ -calculus is basically composed of terms and

commands. We add contexts to the $\lambda\mu$ -calculus. It eases mappings between the $\lambda\mu$ and $\bar{\lambda}\mu\tilde{\mu}$ -calculi. We extend the translation from the $\lambda\mu$ -calculus to the $\bar{\lambda}\mu\tilde{\mu}$ -calculus defined in [Curien and Herbelin, 2000] over the $\lambda\mu$ -contexts. We define backwards a translation from the $\bar{\lambda}\mu\tilde{\mu}$ -calculus to the $\lambda\mu$ -calculus.

In section 2 we present the $\lambda\mu$ -calculus. In section 3 we present the $\bar{\lambda}\mu\tilde{\mu}$ -calculus. In section 4 we define translations between these two calculi. In section 5 we prove simulation theorems that hold for call-by-name and call-by-value.

2 $\lambda\mu$ -calculus

We follow the definition given in [Parigot, 1992]. Firstly, we present the grammar of terms and commands. Secondly, we present the system of simple types. Thirdly, we present generic reductions and their call-by-name and call-by-value projections. Fourthly, we extend both the grammar and the type system to the contexts.

Basically, the $\lambda\mu$ -calculus is composed of terms and commands. They are defined by mutual induction:

$$t ::= x \mid \lambda x.t \mid (t)t \mid \mu\alpha.c \quad c ::= [\alpha]t$$

Symbols x range over λ -variables, symbols α range over μ -variables. We note $x \in t$ or $\alpha \in t$ the fact that x or α has a free occurrence in t . Symbols λ and μ are binders. Two terms are equal modulo α -equivalence.

The system of simple types for the $\lambda\mu$ -calculus is based on two kinds of sequents. The first $\Gamma \vdash t : T \mid \Delta$ concerns the terms and the second $c : (\Gamma \vdash \Delta)$ concerns the commands in which T is a simple type obtained by the grammar $T ::= X \mid T \rightarrow T$, Γ is a finite domain application from λ -variables to simple types and Δ is a finite domain application from μ -variables to simple types. Γ, Γ' denotes the union of the applications Γ and Γ' . System rules are:

$$\begin{array}{c} \frac{}{x:A \vdash x:A \mid} \quad \frac{\Gamma \vdash t:B \mid \Delta}{\Gamma \setminus \{x:A\} \vdash \lambda x.t:A \rightarrow B \mid \Delta} \quad \frac{\Gamma \vdash u:A \rightarrow B \mid \Delta \quad \Gamma' \vdash v:A \mid \Delta'}{\Gamma, \Gamma' \vdash (u) v:B \mid \Delta, \Delta'} (*) \\[10pt] \frac{\Gamma \vdash t:A \mid \Delta}{[\alpha]t:(\Gamma \vdash \Delta, \alpha:A)} (*) \quad \frac{c:(\Gamma \vdash \Delta)}{\Gamma \vdash \mu\alpha.c:A \mid \Delta \setminus \{\alpha:A\}} \end{array}$$

The restriction $(*)$ requires that Γ and Γ' match each other on the intersection of their domains. This holds for Δ and Δ' too.

The category of contexts is introduced in order to ease comparisons with the homonymous category of the $\bar{\lambda}\mu\tilde{\mu}$ -calculus. $\lambda\mu$ -contexts are defined by mutual induction with the terms:

$$e ::= \alpha \mid \beta(t) \mid t \cdot e$$

We can see contexts as commands with a hole to fill. The first construction α expects a term t in order to provide the command $[\alpha]t$. The second $\beta(t)$ expects a term u in order to provide the command $[\beta](t)u$. The last $t \cdot h$ puts the term t on a stack and expects another term to fill the hole.

Definition 1. Let t a term and e a context. The command $e\{t\}$ is defined by induction on e :

$$e\{t\} = \begin{cases} [\alpha] t & \text{if } e = \alpha \\ [\beta](u) t & \text{if } e = \beta(u) \\ h\{(t) u\} & \text{if } e = u \cdot h \end{cases}$$

The type system is extended to another kind of sequents $\Gamma \mid e : T \vdash \Delta$. The typing rules give the context e the type of the term t that fills the hole of e :

$$\frac{}{\mid \alpha : A \vdash \alpha : A} \quad \frac{\Gamma \vdash t : (A \rightarrow B) \mid \Delta}{\Gamma \mid \beta(t) : A \vdash \Delta, \beta : B} \quad \frac{\Gamma \vdash t : A \mid \Delta \quad \Gamma' \mid e : B \vdash \Delta'}{\Gamma, \Gamma' \mid t : e : (A \rightarrow B) \vdash \Delta, \Delta'}$$

A sequent calculus like cut-rule can then be derived in this system as a term against context application.

Lemma 1. The rule $\frac{\Gamma \vdash t : A \mid \Delta \quad \Gamma' \mid e : A \vdash \Delta'}{e\{t\} : (\Gamma, \Gamma' \vdash \Delta, \Delta')}$ holds in $\lambda\mu$.

Proof. By induction on e .

- if $e = \alpha$ then $e\{t\} = [\alpha] t$ and $\frac{\Gamma \vdash t : A \mid \Delta}{[\alpha] t : (\Gamma \vdash \Delta, \alpha : A)}$
- if $e = \beta(u)$ then $e\{t\} = [\beta](u) t$ and

$$\frac{\frac{\Gamma \vdash u : (A \rightarrow B) \mid \Delta \quad \Gamma' \vdash t : A \mid \Delta'}{\Gamma, \Gamma' \vdash (u) t : B \mid \Delta, \Delta'}}{[\beta](u) t : (\Gamma, \Gamma' \vdash \Delta, \Delta', \beta : B)}$$

- if $e = u \cdot h$ then $e\{t\} = h\{(t) u\}$ and

$$\frac{\frac{\Gamma \vdash t : (A \rightarrow B) \mid \Delta \quad \Gamma' \vdash u : A \mid \Delta'}{\Gamma, \Gamma' \vdash (t) u : B \mid \Delta, \Delta'} \quad \Gamma'' \mid h : B \vdash \Delta''}{h\{(t) u\} : (\Gamma, \Gamma', \Gamma'' \vdash \Delta, \Delta', \Delta'')} \text{ind. hyp.}$$

Definition 2. Let t a term, e a context and α a μ -variable, The term $t[\alpha \leftarrow e]$ – the substitution of α by e in t – is defined by induction on t :

$$t[\alpha \leftarrow e] = \begin{cases} x & \text{if } t = x \\ \lambda x. u[\alpha \leftarrow e] & \text{if } t = \lambda x. u \\ (u[\alpha \leftarrow e]) v[\alpha \leftarrow e] & \text{if } t = (u) v \\ \mu \beta. c[\alpha \leftarrow e] & \text{if } t = \mu \beta. c \end{cases}$$

$$c[\alpha \leftarrow e] = \begin{cases} e\{t[\alpha \leftarrow e]\} & \text{if } c = [\alpha] t \\ [\beta] t[\alpha \leftarrow e] & \text{if } c = [\beta] t \end{cases}$$

The computation notion is based on reductions. We remind one-step reduction rules:

$$\begin{array}{ll} (\lambda x. u) t \rightarrow_{\beta} u[x \leftarrow t] & (\mu \alpha. c) t \rightarrow_{\mu} \mu \alpha. c[\alpha \leftarrow t \cdot \alpha] \\ \mu \delta. [\delta] t \rightarrow_{\theta} t \text{ (if } \delta \notin t) & (t) \mu \alpha. c \rightarrow_{\mu'} \mu \alpha. c[\alpha \leftarrow \alpha(t)] \\ & [\beta] \mu \alpha. c \rightarrow_{\rho} c[\alpha \leftarrow \beta] \end{array}$$

The reduction $\xrightarrow{*}_\gamma$ stands for the reflexive and transitive closure of $\rightarrow_\gamma$ and the reduction $\xrightarrow{*}$ stands for the union of $\xrightarrow{*}_\gamma$ for $\gamma \in \{\beta, \mu, \mu', \rho, \theta\}$.

Some of these reductions are linear. Both of the ρ and θ -reductions are linear because they correspond to the identity in NK. The β -reduction from the term $(\lambda x.t)y$ is linear because it consists of replacing a variable by another variable inside a term. It corresponds to a normalisation against an axiom rule in NK. The β -reduction from the term $(\lambda x.t)u$ where x has a single free occurrence in t is linear too because it consists either of substituting a single variable occurrence by any term. It corresponds either to a normalisation without a proof-tree branch duplication.

Reductions $\leadsto_\gamma$, $\xrightarrow{*}_\gamma$ and $\xrightarrow{*}$ have the same meanings as in the general case. The relation $\approx$ is defined as the reflexive, transitive and symmetric closure of $\xrightarrow{*}$.

There exists a critical pair for computation determinism. Applicative terms $(\lambda x.t)\mu\beta.d$ and $(\mu\alpha.c)\mu\beta.d$ can be β or μ' -rewritten in the first case and μ or μ' -rewritten in the second case. We can see the call-by-name and call-by-value disciplines as restrictions of the generic reductions.

The call-by-name evaluation consists of allowing every reduction but the μ' -rule. The β -reduction holds in the first case and the μ -reduction in the second. Formally the call-by-name reduction is $\xrightarrow{*}_n = \xrightarrow{*} \setminus \xrightarrow{*}_{\mu'}$.

The call-by-value evaluation consists of prohibiting β and μ -reductions in which the argument is a μ -abstraction. Formally we define a subset of terms called values by this grammar: $v ::= x \mid \lambda x.t$. β_v and μ_v -reductions are defined instead of generic β and μ ones:

$$(\lambda x.u)v \rightarrow_{\beta_v} u[x \leftarrow v] \quad (\mu\alpha.c)v \rightarrow_{\mu_v} \mu\alpha.c[\alpha \leftarrow v \cdot \alpha]$$

The call-by-value reduction $\xrightarrow{*}_v$ is the union of $\xrightarrow{*}_\gamma$ for $\gamma \in \{\beta_v, \mu_v, \mu', \rho, \theta\}$. Critical pairs are then μ' -rewritten.

There is another way to define call-by-value into the $\lambda\mu$ -calculus. The solution is detailed in [Ong and Stewart, 1997]. It consists of restricting the μ' -rule to values instead of the μ :

$$(v)\mu\alpha.c \rightarrow_{\mu'_v} \mu\alpha.c[\alpha \leftarrow \alpha(v)]$$

Formally $\xrightarrow{*}_v$ becomes the union of $\xrightarrow{*}_\gamma$ for $\gamma \in \{\beta_v, \mu, \mu'_v, \rho, \theta\}$. In fact terms $(\lambda x.t)\mu\alpha.c$ and $(\mu\alpha.c)\mu\alpha'.c'$ are respectively μ' and μ -reduced because $\mu\alpha.c$ is not a value in these cases. However, we follow Curien-Herbelin's call-by-value definition.

We finish this section by a lemma. It is useful for the section 5 simulation theorems. Any command of the form $e\{\mu\alpha.c\}$ is a redex. However, some can not be reduced in call-by-name nor in call-by-value.

Lemma 2. $e\{\mu\alpha.c\} \xrightarrow{*} c[\alpha \leftarrow e]$

Proof. By induction on e .

- if $e = \beta$ then $e\{\mu\alpha.c\} = [\beta]\mu\alpha.c \rightsquigarrow_\rho c[\alpha \leftarrow \beta]$
- if $e = \beta(t)$ then

$$\begin{aligned} e\{\mu\alpha.c\} &= [\beta](t)\mu\alpha.c \\ &\rightarrow_{\mu'} [\beta]\mu\alpha.c[\alpha \leftarrow \alpha(t)] \\ &\rightsquigarrow_\rho c[\alpha \leftarrow \alpha(t)][\alpha \leftarrow \beta] \\ &= c[\alpha \leftarrow \beta(t)] \end{aligned}$$

- if $e = t \cdot h$ then

$$\begin{aligned} e\{\mu\alpha.c\} &= h\{(\mu\alpha.c)t\} \\ &\rightarrow_\mu h\{\mu\alpha.c[\alpha \leftarrow t \cdot \alpha]\} \\ &\xrightarrow{*} c[\alpha \leftarrow t \cdot \alpha][\alpha \leftarrow h] \\ &= c[\alpha \leftarrow t \cdot h] \end{aligned}$$

This lemma does not hold in call-by-name for the $\beta(t)$ induction case because no μ' -rule is allowed. It holds in call-by-value if t is a value for the $h \cdot t$ induction case.

3 $\bar{\lambda}\mu\tilde{\mu}$ -calculus

The $\bar{\lambda}\mu\tilde{\mu}$ -calculus has the same relation against **LK** as the $\lambda\mu$ -calculus against **NK**. Reductions of $\bar{\lambda}\mu\tilde{\mu}$ -calculus correspond to the cut elimination steps in **LK** as well as the $\lambda\mu$ -reductions correspond to the **NK**-normalisation. We follow the definition given in [Curien and Herbelin, 2000]. Firstly, we present the grammar of the $\bar{\lambda}\mu\tilde{\mu}$ -calculus. Secondly, we present the simple type system. Thirdly, we present generic reductions and their call-by-name and call-by-value projections.

The $\bar{\lambda}\mu\tilde{\mu}$ -calculus is basically composed of terms, commands and contexts¹. They are defined by mutual induction:

$$t ::= x \mid \lambda x.t \mid \mu\alpha.c \quad c ::= \langle t \mid e \rangle \quad e ::= \alpha \mid t \cdot e \mid \tilde{\mu}x.c$$

As in the $\lambda\mu$, symbols x range over λ -variables, symbols α range over μ -variables and symbols λ , μ and $\tilde{\mu}$ are binders. Terms are equal modulo α -equivalence.

This calculus symmetry looks like **LK**'s left/right symmetry. It is confirmed by its system of simple types. This system shares types with the $\lambda\mu$ -calculus. It shares the same kinds of sequents too. Its rules are:

$$\begin{array}{c} \frac{}{x:A \vdash x:A \mid} \quad \frac{}{\mid \alpha:A \vdash \alpha:A} \quad \frac{\Gamma \vdash t:B \mid \Delta}{\Gamma \setminus \{x:A\} \vdash \lambda x.t:A \rightarrow B \mid \Delta} \quad \frac{\Gamma \vdash t:A \mid \Delta \quad \Gamma' \mid e:B \vdash \Delta'}{\Gamma, \Gamma' \mid t \cdot e:A \rightarrow B \vdash \Delta, \Delta'} (*) \\[10pt] \frac{c:(\Gamma \vdash \Delta)}{\Gamma \vdash \mu\alpha.c:A \mid \Delta \setminus \{\alpha:A\}} \quad \frac{c:(\Gamma \vdash \Delta)}{\Gamma \setminus \{x:A\} \mid \tilde{\mu}x.c:A \vdash \Delta} \quad \frac{\Gamma \vdash t:A \mid \Delta \quad \Gamma' \mid e:A \vdash \Delta'}{\langle t \mid e \rangle : (\Gamma, \Gamma' \vdash \Delta, \Delta')} (*) \end{array}$$

¹ In [Dougherty et al., 2004] these are referred to respectively callers, callees and capsules. We kept the terminology in [Curien and Herbelin, 2000] that sounds closer to its meaning: terms are programs, contexts are environments and commands represent "a closed system containing both the program and its environment".

The restriction $(*)$ is the same as that of $\lambda\mu$.

We present one-step reduction rules. Substitutions inside the $\bar{\lambda}\mu\tilde{\mu}$ -calculus are supposed to be known. Each rule concerns a command but the θ -rule:

$$\begin{array}{ll} \langle \lambda x.u \mid t \cdot e \rangle \rightarrow_{\beta} \langle t \mid \tilde{\mu}x.\langle u \mid e \rangle \rangle & \langle \mu\alpha.c \mid e \rangle \rightarrow_{\mu} c[\alpha \leftarrow e] \\ \mu\delta.\langle t \mid \delta \rangle \rightarrow_{\theta} t \ (\delta \notin t) & \langle t \mid \tilde{\mu}x.c \rangle \rightarrow_{\tilde{\mu}} c[x \leftarrow t] \end{array}$$

μ and $\tilde{\mu}$ -reductions are duals of each other. They correspond to the structural rules in $\mathbf{LK}$. Reductions $\xrightarrow{*}_{\gamma}$ and $\xrightarrow{\sim}_{\gamma}$ have the same meanings as in the $\lambda\mu$ -calculus. The β -rule is a mere term modification without term duplication. It is therefore a linear reduction. The θ -reduction is linear too. There is no ρ -reduction. It is a μ -rule particular case in which $e = \beta$.

This system is not deterministic. There is a single critical pair $\langle \mu\alpha.c \mid \tilde{\mu}x.d \rangle$. It can be both μ or $\tilde{\mu}$ -rewritten so that Church-Rosser's property does not hold. In fact $\langle \mu\alpha.\langle x \mid y \cdot \alpha \rangle \mid \tilde{\mu}x.\langle z \mid x \cdot \beta \rangle \rangle$ is μ -rewritten as $\langle x \mid y \cdot \tilde{\mu}x.\langle z \mid x \cdot \beta \rangle \rangle$ and is $\tilde{\mu}$ -rewritten as $\langle z \mid \mu\alpha.\langle x \mid y \cdot \alpha \rangle \cdot \beta \rangle$. These are two different normal forms.

Call-by-name and call-by-value disciplines still deal with this problem. They both consist of restricting the context construction. The first new grammar is called $\bar{\lambda}\mu\tilde{\mu}_T$ and the second is called $\bar{\lambda}\mu\tilde{\mu}_Q$.

The call-by-name evaluation consists of restricting the μ -rule to a subset of contexts that are called stacks. $\bar{\lambda}\mu\tilde{\mu}_T$ -grammar is:

$$t ::= x \mid \lambda x.t \mid \mu\alpha.c \quad c ::= \langle t \mid e \rangle \quad s ::= \alpha \mid t \cdot s \quad e ::= s \mid \tilde{\mu}x.c$$

The μ_n -rule is restricted to the stacks:

$$\langle \mu\alpha.c \mid s \rangle \rightarrow_{\mu_n} c[\alpha \leftarrow s]$$

Call-by-name reduction $\xrightarrow{*}_n$ is the union of $\xrightarrow{*}_{\gamma}$ for $\gamma \in \{\beta, \mu_n, \tilde{\mu}, \theta\}$. The critical pair can then only be $\tilde{\mu}$ -rewritten. This reduction was proved confluent and stable in the $\bar{\lambda}\mu\tilde{\mu}_T$ -calculus in [Curien and Herbelin, 2000].

The call-by-value oriented grammar consists of allowing the $t \cdot e$ context construction only for values. $\bar{\lambda}\mu\tilde{\mu}_Q$ -grammar is:

$$t ::= x \mid \lambda x.t \mid \mu\alpha.c \quad v ::= x \mid \lambda x.t \quad c ::= \langle t \mid e \rangle \quad e ::= \alpha \mid v \cdot e \mid \tilde{\mu}x.c$$

The $\tilde{\mu}_v$ -rule is restricted to values:

$$\langle v \mid \tilde{\mu}x.c \rangle \rightarrow_{\tilde{\mu}_v} c[x \leftarrow v]$$

Call-by-value reduction $\xrightarrow{*}_v$ is the union of $\xrightarrow{*}_{\gamma}$ for $\gamma \in \{\beta, \mu, \tilde{\mu}_v, \theta\}$. The command $\langle \mu\alpha.c \mid \mu\alpha'.c' \rangle$ can then only be μ -rewritten. This reduction was proved confluent and stable in the $\bar{\lambda}\mu\tilde{\mu}_Q$ -calculus in [Curien and Herbelin, 2000].

The β' -rule contracts as shortcut for both a linear β -rule and a $\tilde{\mu}$ -rule:

$$\langle \lambda x.u \mid t \cdot e \rangle \rightarrow_{\beta'} \langle u[x \leftarrow t] \mid e \rangle$$

This β' -rule is obviously compatible with the call-by-name evaluation. It is also compatible with the call-by-value because t is a value by definition of $\bar{\lambda}\mu\tilde{\mu}_Q$.

4 Translations between $\lambda\mu$ and $\bar{\lambda}\mu\tilde{\mu}$ -calculi

We define a translation $(\)^\dagger$ from $\lambda\mu$ to $\bar{\lambda}\mu\tilde{\mu}$. It extends that of Curien-Herbelin to the $\lambda\mu$ -contexts. We define backwards a translation $(\)^\circ$ from $\bar{\lambda}\mu\tilde{\mu}$ to $\lambda\mu$. We prove properties about their compatibilities with the simple type system and about their compositions.

Definition 3. Application $(\)^\dagger$ maps any $\lambda\mu$ -term t , command c and context e respectively to a $\bar{\lambda}\mu\tilde{\mu}$ -term, command and context. $(\)^\dagger$ is defined by induction on t , c and e :

$$t^\dagger = \begin{cases} x & \text{if } t = x \\ \lambda x.u^\dagger & \text{if } t = \lambda x.u \\ \mu\beta.\langle v^\dagger \mid \tilde{\mu}y.\langle u^\dagger \mid y \cdot \beta \rangle \rangle & \text{if } t = (u)v \quad (\star) \\ \mu\alpha.c^\dagger & \text{if } t = \mu\alpha.c \end{cases}$$

$$c^\dagger = [\alpha] t^\dagger = \langle t^\dagger | \alpha \rangle$$

$$e^\dagger = \begin{cases} \alpha & \text{if } e = \alpha \\ \tilde{\mu}y.\langle t^\dagger \mid y \cdot \beta \rangle & \text{if } e = \beta(t) \\ t^\dagger \cdot h^\dagger & \text{if } e = t \cdot h \end{cases} \quad (\star\star)$$

Condition $(\star)$ requires that variables y and β have no free occurrence in u neither in v . Condition $(\star\star)$ requires that $y \notin t$. A straightforward induction leads us to state that t and $t^\dagger$ have the same free variables set.

It seems more natural to translate $(u) v$ by $\mu\beta.\langle u^\dagger | v^\dagger \cdot \beta \rangle$. This shorter term corresponds in LK to the arrow elimination rule in NK too. But it would not be compatible with the call-by-value evaluation. For example, $(x) \mu\alpha.c$ would be translated as $\mu\beta.\langle x | \mu\alpha.c^\dagger \cdot \beta \rangle$ in this case. It can not be reduced by any rule in the $\bar{\lambda}\mu\tilde{\mu}$ -calculus. However, $(x) \mu\alpha.c$ can be μ' -reduced in the $\lambda\mu$ -calculus.

$(u)v$ should be translated as $\mu\beta.\langle u^\dagger \mid \tilde{\mu}y.\langle v^\dagger \mid \tilde{\mu}x.\langle y \mid x \cdot \beta \rangle \rangle \rangle$ with Ong and Stewart’s call-by-value definition in [Ong and Stewart, 1997].

We show that translation $(\)^\dagger$ is compatible with the type system. If a typing environment for a term t exists, it holds for $t^\dagger$.

Lemma 3. $\Gamma \vdash t : A \mid \Delta \implies \Gamma \vdash t^\dagger : A \mid \Delta$

Proof. By a straightforward induction on t . We show the less than obvious cases.

- if $t = (u) v$ then $t^\dagger = \mu\beta.\langle v^\dagger \mid \tilde{\mu}y.\langle u^\dagger \mid y \cdot \beta \rangle \rangle$ and

$$\frac{\Gamma' \vdash v^\dagger : A \mid \Delta'}{\frac{\frac{\frac{\Gamma \vdash u^\dagger : A \rightarrow B \mid \Delta}{\frac{y : A \vdash y : A \mid \quad \beta : B \vdash \beta : B}{y : A \mid y : \beta : (A \rightarrow B) \vdash \beta : B}}{\frac{\langle u^\dagger \mid y : \beta \rangle : (\Gamma, y : A \vdash \Delta, \beta : B)}{\Gamma \mid \tilde{\mu} y . \langle u^\dagger \mid y : \beta \rangle \vdash \Delta, \beta : B}}}{\frac{\langle v^\dagger \mid \tilde{\mu} y . \langle u^\dagger \mid y : \beta \rangle \rangle : (\Gamma, \Gamma' \vdash \Delta, \Delta', \beta : B)}{\Gamma, \Gamma' \vdash \mu \beta . \langle v^\dagger \mid \tilde{\mu} y . \langle u^\dagger \mid y : \beta \rangle \rangle : B \mid \Delta, \Delta'}}$$

– if $e = \beta(t)$ then $e^\dagger = \tilde{\mu}y.\langle t^\dagger \mid y \cdot \beta \rangle$ and

$$\frac{\frac{\Gamma \vdash t^\dagger : (A \rightarrow B) \mid \Delta \quad \frac{\frac{y:A \vdash y:A \mid \quad \mid \beta:B \vdash \beta:B}{y:A \mid y:\beta:(A \rightarrow B) \vdash \beta:B}}{\langle t^\dagger \mid y \cdot \beta \rangle : (\Gamma, y:A \vdash \Delta, \beta:B)}}{\Gamma \mid \tilde{\mu}y.\langle t^\dagger \mid y \cdot \beta \rangle : A \vdash \Delta, \beta:B}$$

Definition 4. Application $()^\circ$ maps backwards any $\bar{\lambda}\mu\tilde{\mu}$ -term t to a $\lambda\mu$ -term. Definition 1 is used to translate any $\bar{\lambda}\mu\tilde{\mu}$ -command c . Definition of the $\lambda\mu$ -contexts is used to map the $\bar{\lambda}\mu\tilde{\mu}$ -contexts e as well. $()^\circ$ is built by induction on t , c and e :

$$t^\circ = \begin{cases} x & \text{if } t = x \\ \lambda x.u^\circ & \text{if } t = \lambda x.u \\ \mu\alpha.c^\circ & \text{if } t = \mu\alpha.c \end{cases}$$

$$c^\circ = \langle t \mid e \rangle^\circ = e^\circ \{t^\circ\}$$

$$e^\circ = \begin{cases} \alpha & \text{if } e = \alpha \\ t^\circ \cdot h^\circ & \text{if } e = t \cdot h \\ \beta(\lambda x.\mu\delta.c^\circ) & \text{if } e = \tilde{\mu}x.c \quad (*) \end{cases}$$

Condition $(*)$ requires that $\delta \notin c$. t and t° have the same free variables set. Application $()^\circ$ is compatible with the type system too.

Lemma 4. $\Gamma \vdash t : A \mid \Delta \implies \Gamma \vdash t^\circ : A \mid \Delta$

Proof. By a straightforward induction on t . We give two cases.

– if $c = \langle t \mid e \rangle$ then $c^\circ = e^\circ \{t^\circ\}$ and

$$\frac{\frac{\Gamma \vdash t:A \mid \Delta \quad \Gamma' \mid e:A \vdash \Delta'}{(t \mid e) : (\Gamma, \Gamma' \vdash \Delta, \Delta')}}{\Gamma \vdash t^\circ : A \mid \Delta \quad \Gamma' \mid e^\circ : A \vdash \Delta'} \text{lem. 1}$$

– if $e = \tilde{\mu}x.c$ then $e^\circ = \beta(\lambda x.\mu\delta.c^\circ)$ and

$$\frac{c : (\Gamma \vdash \Delta)}{\Gamma \setminus \{x:A\} \mid \tilde{\mu}x.c : A \vdash \Delta}^\circ = \frac{\frac{c^\circ : (\Gamma \vdash \Delta)}{\Gamma \vdash \mu\delta.c^\circ : B \mid \Delta}}{\Gamma \setminus \{x:A\} \vdash \lambda x.\mu\delta.c^\circ : (A \rightarrow B)} \quad \Gamma \setminus \{x:A\} \mid \beta(\lambda x.\mu\delta.c^\circ) : A \vdash \Delta, \beta:B$$

We focus on properties about the composition of $()^\dagger$ and $()^\circ$. We want to state that $t^{\dagger\circ} = t$ and that $t^{\circ\dagger} = t$ for any term. But it is not the case, these results hold modulo linear reductions.

Theorem 1. $t^{\dagger\circ} \rightsquigarrow^* t$

Proof. By a straightforward induction on t . Every cases is obtained successively by expanding definitions 3, 1, 4 and by applying the induction hypothesis. We give the case which uses linear reductions additionally.

– if $t = (u) v$ then

$$\begin{aligned}
(u) v^{\dagger\circ} &= \mu\beta.\langle v^\dagger | \tilde{\mu}y.\langle u^\dagger | y \cdot \beta \rangle \rangle^\circ \\
&= \mu\beta.[\gamma] (\lambda y.\mu\delta.[\beta] (u^{\dagger\circ}) y) v^{\dagger\circ} \\
&\stackrel{*}{\rightsquigarrow} \mu\beta.[\gamma] (\lambda y.\mu\delta.[\beta] (u) y) v \\
&\rightsquigarrow_\beta \mu\beta.[\gamma] \mu\delta.[\beta] (u) v \\
&\rightsquigarrow_\rho \mu\beta.[\beta] (u) v \\
&\rightsquigarrow_\theta (u) v
\end{aligned}$$

We prove two lemmas before stating backwards that $()^{\dagger\circ}$ is the identity modulo linear reductions. The first lemma is useful to prove the second.

Lemma 5. $\langle t_0 t_1 \dots t_n^\dagger | e \rangle \stackrel{*}{\rightsquigarrow} \langle t_0^\dagger | t_1^\dagger \cdot \dots \cdot t_n^\dagger \cdot e \rangle$

Proof. By induction on n .

– if $n = 0$ then it is obvious
– if $n = m + 1$ then

$$\begin{aligned}
\langle t_0 t_1 \dots t_m t_{m+1}^\dagger | e \rangle &= \langle \mu\beta.\langle t_{m+1}^\dagger | \tilde{\mu}y.\langle t_0 t_1 \dots t_m^\dagger | y \cdot \beta \rangle \rangle | e \rangle \\
&\rightsquigarrow_\mu \langle t_{m+1}^\dagger | \tilde{\mu}y.\langle t_0 t_1 \dots t_m^\dagger | y \cdot e \rangle \rangle \\
&\rightsquigarrow_{\tilde{\mu}} \langle t_0 t_1 \dots t_m^\dagger | t_{m+1}^\dagger \cdot e \rangle \\
&\stackrel{*}{\rightsquigarrow} \langle t_0^\dagger | t_1^\dagger \cdot \dots \cdot t_m^\dagger \cdot t_{m+1}^\dagger \cdot e \rangle
\end{aligned}$$

The second lemma shows how to map a definition 1 command.

Lemma 6. $e\{t\}^\dagger \stackrel{*}{\rightsquigarrow} \langle t^\dagger | e^\dagger \rangle$

Proof. By induction on e .

– if $e = \alpha$ then it is obvious by definitions 1 and 3
– if $e = \beta(u)$ then

$$\begin{aligned}
\beta(u)\{t\}^\dagger &= [\beta] (u) t^\dagger \\
&= \langle \mu\gamma.\langle t^\dagger | \tilde{\mu}y.\langle u^\dagger | y \cdot \gamma \rangle \rangle | \beta \rangle \\
&\rightsquigarrow_\mu \langle t^\dagger | \tilde{\mu}y.\langle u^\dagger | y \cdot \beta \rangle \rangle \\
&= \langle t^\dagger | \beta(u)^\dagger \rangle
\end{aligned}$$

– if $e = u \cdot h$ then

$$\begin{aligned}
u \cdot h\{t\}^\dagger &= h\{t\} u^\dagger \\
&\stackrel{*}{\rightsquigarrow} \langle (t) u^\dagger | h^\dagger \rangle \\
&\stackrel{*}{\rightsquigarrow} \langle t^\dagger | u^\dagger \cdot h^\dagger \rangle \\
&= \langle t^\dagger | u \cdot h^\dagger \rangle
\end{aligned}$$

Theorem 2. $t^{\circ\dagger} \stackrel{*}{\rightsquigarrow} t$

Proof. By induction on t . We apply definitions 3, 4 successively and the induction hypothesis. We give a typical case and another which needs either the previous lemma or linear reductions.

– if $c = \langle t \mid e \rangle$ then

$$\begin{aligned} \langle t \mid e \rangle^{\circ\dagger} &= e^{\circ} \{t^{\circ}\}^{\dagger} \\ &\stackrel{*}{\rightsquigarrow} \langle t^{\circ\dagger} \mid e^{\circ\dagger} \rangle \\ &\stackrel{*}{\rightsquigarrow} \langle t \mid e \rangle \end{aligned}$$

– if $e = \tilde{\mu}x.c$ then

$$\begin{aligned} \tilde{\mu}x.c^{\circ\dagger} &= \beta(\lambda x.\mu\beta.c^{\circ})^{\dagger} \\ &= \tilde{\mu}y.\langle \lambda x.\mu\beta.c^{\circ\dagger} \mid y \cdot \beta \rangle \\ &\stackrel{*}{\rightsquigarrow} \tilde{\mu}y.\langle \lambda x.\mu\beta.c \mid y \cdot \beta \rangle \\ &\rightsquigarrow_{\beta} \tilde{\mu}y.\langle y \mid \tilde{\mu}x.\langle \mu\beta.c \mid \beta \rangle \rangle \\ &\rightsquigarrow_{\tilde{\mu}} \tilde{\mu}x.\langle \mu\beta.c \mid \beta \rangle \\ &\rightsquigarrow_{\mu} \tilde{\mu}x.c \end{aligned}$$

5 Simulations between $\lambda\mu$ and $\bar{\lambda}\mu\tilde{\mu}$ -calculi

We want to prove that the $\lambda\mu$ -calculus simulates and is simulated backwards by the $\bar{\lambda}\mu\tilde{\mu}$ -calculus. We focus on the undirected evaluation. Call-by-name and call-by-value are drawn from this.

We begin with the simulation of the $\lambda\mu$ by the $\bar{\lambda}\mu\tilde{\mu}$. The next four lemmas show results of a $\lambda\mu$ -substitution after a β , μ , μ' and ρ -reduction. Each proof consists successively of

- expanding the $\lambda\mu$ -substitution
- expanding the definition of $()^{\dagger}$
- applying the induction hypothesis if necessary
- factorising the $\bar{\lambda}\mu\tilde{\mu}$ -substitution
- factorising the definition of $()^{\dagger}$

We give basic cases and those which use lemmas additionally for any proof.

Lemma 7. $t[x \leftarrow u]^{\dagger} = t^{\dagger}[x \leftarrow u^{\dagger}]$

Proof. By induction on t .

- if $t = x$ then $x[x \leftarrow u]^{\dagger} = u^{\dagger} = x^{\dagger}[x \leftarrow u^{\dagger}]$
- if $t = y$ then $y[x \leftarrow u]^{\dagger} = y = y^{\dagger}[x \leftarrow u^{\dagger}]$
- if $t = (v)w$ then

$$\begin{aligned} (v)w[x \leftarrow u]^{\dagger} &= (v[x \leftarrow u])w[x \leftarrow u]^{\dagger} \\ &= \mu\beta.\langle w[x \leftarrow u]^{\dagger} \mid \tilde{\mu}y.\langle v[x \leftarrow u]^{\dagger} \mid y \cdot \beta \rangle \rangle \\ &= \mu\beta.\langle w^{\dagger}[x \leftarrow u^{\dagger}] \mid \tilde{\mu}y.\langle v^{\dagger}[x \leftarrow u^{\dagger}] \mid y \cdot \beta \rangle \rangle \\ &= \mu\beta.\langle w^{\dagger} \mid \tilde{\mu}y.\langle v^{\dagger} \mid y \cdot \beta \rangle \rangle [x \leftarrow u^{\dagger}] \\ &= (v)w^{\dagger}[x \leftarrow u^{\dagger}] \end{aligned}$$

Lemma 8. $t[\alpha \leftarrow u \cdot \alpha]^{\dagger} \stackrel{*}{\rightsquigarrow} t^{\dagger}[\alpha \leftarrow u^{\dagger} \cdot \alpha]$

Proof. By induction on t .

– if $t = (a) b$ then

$$\begin{aligned}
(a) b[\alpha \leftarrow u \cdot \alpha]^\dagger &= (a[\alpha \leftarrow u \cdot \alpha]) b[\alpha \leftarrow u \cdot \alpha]^\dagger \\
&= \mu\beta. \langle b[\alpha \leftarrow u \cdot \alpha]^\dagger | \tilde{\mu}y. \langle a[\alpha \leftarrow u \cdot \alpha]^\dagger | y \cdot \beta \rangle \rangle \\
&\stackrel{*}{\rightsquigarrow} \mu\beta. \langle b^\dagger[\alpha \leftarrow u^\dagger \cdot \alpha] | \tilde{\mu}y. \langle a^\dagger[\alpha \leftarrow u^\dagger \cdot \alpha] | y \cdot \beta \rangle \rangle \\
&= \mu\beta. \langle b^\dagger | \tilde{\mu}y. \langle a^\dagger | y \cdot \beta \rangle \rangle [\alpha \leftarrow u^\dagger \cdot \alpha] \\
&= (a) b^\dagger[\alpha \leftarrow u^\dagger \cdot \alpha]
\end{aligned}$$

– if $c = [\alpha] w$ then

$$\begin{aligned}
[\alpha] w[\alpha \leftarrow u \cdot \alpha]^\dagger &= [\alpha] w[\alpha \leftarrow u \cdot \alpha]^\dagger \\
&= u \cdot \alpha \{w[\alpha \leftarrow u \cdot \alpha]\}^\dagger \\
&= \langle w[\alpha \leftarrow u \cdot \alpha]^\dagger | u^\dagger \cdot \alpha \rangle \\
&\stackrel{*}{\rightsquigarrow} \langle w^\dagger[\alpha \leftarrow u^\dagger \cdot \alpha] | u^\dagger \cdot \alpha \rangle \\
&= \langle w^\dagger | \alpha \rangle [\alpha \leftarrow u^\dagger \cdot \alpha] \\
&= [\alpha] w^\dagger[\alpha \leftarrow u^\dagger \cdot \alpha]
\end{aligned}$$

Lemma 9. $t[\alpha \leftarrow \alpha(u)]^\dagger \stackrel{*}{\rightsquigarrow} t^\dagger[\alpha \leftarrow \tilde{\mu}y. \langle y | y \cdot \alpha \rangle] u^\dagger$

Proof. By induction on t .

– if $t = (a) b$ then

$$\begin{aligned}
(a) b[\alpha \leftarrow \alpha(u)]^\dagger &= (a[\alpha \leftarrow \alpha(u)]) b[\alpha \leftarrow \alpha(u)]^\dagger \\
&= \mu\beta. \langle b[\alpha \leftarrow \alpha(u)]^\dagger | \tilde{\mu}y. \langle a[\alpha \leftarrow \alpha(u)]^\dagger | y \cdot \beta \rangle \rangle \\
&\stackrel{*}{\rightsquigarrow} \mu\beta. \langle b^\dagger[\alpha \leftarrow \tilde{\mu}y. \langle u^\dagger | y \cdot \alpha \rangle] | \tilde{\mu}y. \langle a^\dagger[\alpha \leftarrow \tilde{\mu}y. \langle u^\dagger | y \cdot \alpha \rangle] | y \cdot \beta \rangle \rangle \\
&= \mu\beta. \langle b^\dagger | \tilde{\mu}y. \langle a^\dagger | y \cdot \beta \rangle \rangle [\alpha \leftarrow \tilde{\mu}y. \langle u^\dagger | y \cdot \alpha \rangle] \\
&= (a) b^\dagger[\alpha \leftarrow \tilde{\mu}y. \langle u^\dagger | y \cdot \alpha \rangle]
\end{aligned}$$

– if $c = [\alpha] w$ then

$$\begin{aligned}
[\alpha] w[\alpha \leftarrow \alpha(u)]^\dagger &= [\alpha] (w[\alpha \leftarrow \alpha(u)]) u^\dagger \\
&= \langle (w[\alpha \leftarrow \alpha(u)]) u^\dagger | \alpha \rangle \\
&\stackrel{*}{\rightsquigarrow} \langle w[\alpha \leftarrow \alpha(u)]^\dagger | u^\dagger \cdot \alpha \rangle \\
&\stackrel{*}{\rightsquigarrow} \langle w^\dagger[\alpha \leftarrow \tilde{\mu}y. \langle u^\dagger | y \cdot \alpha \rangle] | u^\dagger \cdot \alpha \rangle \\
&= \langle w^\dagger | \alpha \rangle [\alpha \leftarrow \tilde{\mu}y. \langle u^\dagger | y \cdot \alpha \rangle] \\
&= [\alpha] w^\dagger[\alpha \leftarrow \tilde{\mu}y. \langle u^\dagger | y \cdot \alpha \rangle]
\end{aligned}$$

Lemma 10. $t[\alpha \leftarrow \beta]^\dagger = t^\dagger[\alpha \leftarrow \beta]$

Proof. By induction on t .

– if $c = [\alpha] u$ then

$$\begin{aligned}
[\alpha] u[\alpha \leftarrow \beta]^\dagger &= [\beta] u[\alpha \leftarrow \beta]^\dagger \\
&= \langle u[\alpha \leftarrow \beta]^\dagger | \beta \rangle \\
&= \langle u^\dagger[\alpha \leftarrow \beta] | \beta \rangle \\
&= \langle u^\dagger | \alpha \rangle [\alpha \leftarrow \beta] \\
&= [\alpha] u^\dagger[\alpha \leftarrow \beta]
\end{aligned}$$

Theorem 3 (simulation of the $\lambda\mu$ -calculus by the $\bar{\lambda}\mu\tilde{\mu}$ -calculus).

$$t \rightarrow_\gamma v \implies \exists u \, t^\dagger \xrightarrow{*} u \wedge v^\dagger \xrightarrow{*} u$$

Proof. By cases on γ .

– if $\gamma = \beta$ then

$$\begin{aligned} (\lambda x.u) v^\dagger &= \mu\beta.\langle v^\dagger \mid \tilde{\mu}y.\langle \lambda x.u^\dagger \mid y \cdot \beta \rangle \rangle \\ &\rightsquigarrow_\beta \mu\beta.\langle v^\dagger \mid \tilde{\mu}y.\langle y \mid \tilde{\mu}x.\langle u^\dagger \mid \beta \rangle \rangle \rangle \\ &\rightsquigarrow_{\tilde{\mu}} \mu\beta.\langle v^\dagger \mid \tilde{\mu}x.\langle u^\dagger \mid \beta \rangle \rangle \\ &\rightarrow_{\tilde{\mu}} \mu\beta.\langle u^\dagger[x \leftarrow v^\dagger] \mid \beta \rangle \\ &\rightsquigarrow_\theta u^\dagger[x \leftarrow v^\dagger] \\ &= u[x \leftarrow v]^\dagger \end{aligned}$$

– if $\gamma = \mu$ then

$$\begin{aligned} (\mu\alpha.c) v^\dagger &= \mu\beta.\langle v^\dagger \mid \tilde{\mu}y.\langle \mu\alpha.c^\dagger \mid y \cdot \beta \rangle \rangle \\ &\rightsquigarrow_{\tilde{\mu}} \mu\alpha.\langle \mu\alpha.c^\dagger \mid v^\dagger \cdot \alpha \rangle \\ &\rightarrow_\mu \mu\alpha.c^\dagger[\alpha \leftarrow v^\dagger \cdot \alpha] \\ &\approx \mu\alpha.c[\alpha \leftarrow v \cdot \alpha]^\dagger \end{aligned}$$

– if $\gamma = \mu'$ then

$$\begin{aligned} (v) \mu\alpha.c^\dagger &= \mu\beta.\langle \mu\alpha.c^\dagger \mid \tilde{\mu}y.\langle v^\dagger \mid y \cdot \beta \rangle \rangle \\ &\rightarrow_\mu \mu\alpha.c^\dagger[\alpha \leftarrow \tilde{\mu}y.\langle v^\dagger \mid y \cdot \alpha \rangle] \\ &\approx \mu\alpha.c[\alpha \leftarrow \alpha(v)]^\dagger \end{aligned}$$

– if $\gamma = \rho$ then

$$\begin{aligned} [\beta] \mu\alpha.c^\dagger &= \langle \mu\alpha.c^\dagger \mid \beta \rangle \\ &\rightsquigarrow_\mu c^\dagger[\alpha \leftarrow \beta] \\ &= c[\alpha \leftarrow \beta]^\dagger \end{aligned}$$

– if $\gamma = \theta$ then $\mu\delta.[\delta] t^\dagger = \mu\delta.\langle t^\dagger \mid \delta \rangle \rightsquigarrow_\theta t^\dagger$

Corollary 1 (call-by-name case). $t \rightarrow_n v \implies \exists u \, t^\dagger \xrightarrow{*}_n u \wedge v^\dagger \xrightarrow{*}_n u$

Proof. By cases on β and μ -rules.

$(\lambda x.u) v$ is β -reduced in call-by-name without any restriction. It is simulated in the $\bar{\lambda}\mu\tilde{\mu}$ -calculus by a $\tilde{\mu}$ -reduction. The latter is in call-by-name without any restriction too.

$(\mu\alpha.c) v$ is μ -reduced in call-by-name without any restriction. It is simulated in the $\bar{\lambda}\mu\tilde{\mu}$ -calculus by a μ -reduction. The latter is in call-by-name if $v^\dagger \cdot \alpha$ is a stack. It is the case by definition 3.

Corollary 2 (call-by-value case). $t \rightarrow_v v \implies \exists u \, t^\dagger \xrightarrow{*}_v u \wedge v^\dagger \xrightarrow{*}_v u$

Proof. By cases on β , μ and μ' -rules.

$(\lambda x.u)v$ is β -reduced in call-by-value if v is a value. It is simulated in the $\bar{\lambda}\mu\tilde{\mu}$ -calculus by a $\tilde{\mu}$ -reduction. The latter is in call-by-value if $v^\dagger$ is a value. It is the case by the definition of $\bar{\lambda}\mu\tilde{\mu}_Q$.

$(\mu\alpha.c)v$ is μ -reduced in call-by-value if v is a value. It is simulated in the $\bar{\lambda}\mu\tilde{\mu}$ -calculus by a μ -reduction. The latter is in call-by-value without any restriction.

$(v)\mu\alpha.c$ is μ' -reduced in call-by-value without any restriction. It is simulated in the $\bar{\lambda}\mu\tilde{\mu}$ -calculus by a μ -reduction. The latter is in call-by-value without any restriction as well.

The $\bar{\lambda}\mu\tilde{\mu}$ -simulation by the $\lambda\mu$ -calculus requires preliminary lemmas showing that $(\)^\circ$ commutes over $\lambda\mu$ and $\bar{\lambda}\mu\tilde{\mu}$ -substitutions. Each proof consists of

- expanding the $\bar{\lambda}\mu\tilde{\mu}$ -substitution
- expanding the definition of $(\)^\circ$
- applying the induction hypothesis if necessary
- factorising the $\lambda\mu$ -substitution
- factorising the definition of $(\)^\circ$

Lemma 11. $t[x \leftarrow u]^\circ = t^\circ[x \leftarrow u^\circ]$

Proof. By induction on t .

- if $t = x$ then $x[x \leftarrow u]^\circ = u^\circ = x^\circ[x \leftarrow u^\circ]$
- if $t = y$ then $y[x \leftarrow u]^\circ = y = y^\circ[x \leftarrow u^\circ]$
- if $t = \langle t \mid e \rangle$ then

$$\begin{aligned} \langle t \mid e \rangle[x \leftarrow u]^\circ &= \langle t[x \leftarrow u] \mid e[x \leftarrow u] \rangle^\circ \\ &= e[x \leftarrow u]^\circ \{t[x \leftarrow u]^\circ\} \\ &= e^\circ[x \leftarrow u^\circ] \{t^\circ[x \leftarrow u^\circ]\} \\ &= e^\circ\{t^\circ\}[x \leftarrow u^\circ] \\ &= \langle t \mid e \rangle^\circ[x \leftarrow u^\circ] \end{aligned}$$

Lemma 12. $t[\alpha \leftarrow h]^\circ = t^\circ[\alpha \leftarrow h^\circ]$

Proof. By induction on t .

- if $c = \langle t \mid e \rangle$ then

$$\begin{aligned} \langle t \mid e \rangle[\alpha \leftarrow h]^\circ &= \langle t[\alpha \leftarrow h] \mid e[\alpha \leftarrow h] \rangle^\circ \\ &= e[\alpha \leftarrow h]^\circ \{t[\alpha \leftarrow h]^\circ\} \\ &= e^\circ[\alpha \leftarrow h^\circ] \{t^\circ[\alpha \leftarrow h^\circ]\} \\ &= e^\circ\{t^\circ\}[\alpha \leftarrow h^\circ] \\ &= \langle t \mid e \rangle^\circ[\alpha \leftarrow h^\circ] \end{aligned}$$

- if $e = \alpha$ then $\alpha[\alpha \leftarrow h]^\circ = h^\circ = \alpha^\circ[\alpha \leftarrow h^\circ]$
- if $e = \beta$ then $\beta[\alpha \leftarrow h]^\circ = \beta^\circ = \beta^\circ[\alpha \leftarrow h^\circ]$

Theorem 4 (simulation of the $\bar{\lambda}\mu\tilde{\mu}$ -calculus by the $\lambda\mu$ -calculus).

$$t \rightarrow_\gamma v \implies \exists u \, t^\circ \xrightarrow{*} u \xrightarrow{*} v^\circ$$

Proof. By cases on γ .

– if $\gamma = \beta'$ then

$$\begin{aligned} \langle \lambda x.u \mid v \cdot e \rangle^\circ &= v^\circ \cdot e^\circ \{ \lambda x.u^\circ \} \\ &= e^\circ \{ (\lambda x.u^\circ) v^\circ \} \\ &\rightarrow_\beta e^\circ \{ u^\circ [x \leftarrow v^\circ] \} \\ &= e^\circ \{ u[x \leftarrow v]^\circ \} \\ &= \langle u[x \leftarrow v] \mid e \rangle^\circ \end{aligned}$$

– if $\gamma = \mu$ then

$$\begin{aligned} \langle \mu\alpha.c \mid e \rangle^\circ &= e^\circ \{ \mu\alpha.c^\circ \} \\ &\xrightarrow{*} c^\circ [\alpha \leftarrow e^\circ] \\ &= c[\alpha \leftarrow e]^\circ \end{aligned}$$

– if $\gamma = \tilde{\mu}$ then

$$\begin{aligned} \langle t \mid \tilde{\mu}x.c \rangle^\circ &= [\beta] (\lambda x.\mu\delta.c^\delta) t^\circ \\ &\rightarrow_\beta [\beta] \mu\delta.c^\circ [x \leftarrow t^\circ] \\ &\rightsquigarrow_\rho c^\circ [x \leftarrow t^\circ] \\ &= c[x \leftarrow t]^\circ \end{aligned}$$

– if $\gamma = \theta$ then $\mu\delta.\langle t \mid \delta \rangle^\circ = \mu\delta.[\delta] t^\circ \rightsquigarrow_\theta t^\circ$

Corollary 3 (call-by-name case). $t \rightarrow_n v \implies \exists u \, t^\circ \xrightarrow{*}_n u \xrightarrow{*}_n v^\circ$

Proof. By cases on β' , μ and $\tilde{\mu}$ -rules.

$\langle \lambda x.u \mid v \cdot e \rangle$ is β' -reduced in call-by-name without any restriction. It is simulated in the $\lambda\mu$ -calculus by a β -reduction. The latter is in call-by-name without any restriction too.

$\langle \mu\alpha.c \mid e \rangle$ is μ -reduced in call-by-name if $e \neq \tilde{\mu}x.c'$ else it were $\tilde{\mu}$ -reduced. It is simulated in the $\lambda\mu$ -calculus with the help of lemma 2. The latter is in call-by-name if $e^\circ \neq \beta(t)$ i.e. if $e \neq \tilde{\mu}x.c'$. It is the case by definition 4.

$\langle t \mid \tilde{\mu}x.c \rangle$ is $\tilde{\mu}$ -reduced in call-by-name without any restriction. It is simulated in the $\lambda\mu$ -calculus by a β -reduction. The latter is in call-by-name without any restriction as well.

Corollary 4 (call-by-value case). $t \rightarrow_v v \implies \exists u \, t^\circ \xrightarrow{*}_v u \xrightarrow{*}_v v^\circ$

Proof. By cases on β' , μ and $\tilde{\mu}$ -rules.

$\langle \lambda x.u \mid v \cdot e \rangle$ is β' -reduced in call-by-value if v is a value. It is simulated in the $\lambda\mu$ -calculus by a β -reduction. The latter is in call-by-value if v° is a value. It is the case by definition 4.

$\langle \mu\alpha.c \mid e \rangle$ is μ -reduced in call-by-value if e is either a μ -variable or a context of the form $v \cdot h$ where v is a value or a μ -abstraction by the definition of $\bar{\lambda}\mu\tilde{\mu}_Q$. It is simulated in the $\lambda\mu$ -calculus with the help of lemma 2. The latter is in

call-by-value if v° is a value in a context of the form $h^\circ \cdot v^\circ$ i.e. if v is a value in a $v \cdot h$ context. It is the case by definition 4.

$\langle t \mid \tilde{\mu}x.c \rangle$ is $\tilde{\mu}$ -reduced in call-by-value if t is a value. It is simulated in the $\lambda\mu$ -calculus by a β -reduction. The latter is in call-by-value if t° is a value. It is the case by definition 4.

6 Conclusion

Analysis of the $\lambda\mu$ and $\bar{\lambda}\mu\tilde{\mu}$ -calculi has shown their computational equivalence. It holds for undirected evaluations of pure calculi (see theorems 3 and 4). This result is then easily obtained for call-by-name and call-by-value evaluations (see corollaries 1, 2, 3 and 4). It concerns the simple type system too (see lemmas 3 and 4).

The simulation of the $\bar{\lambda}\mu\tilde{\mu}$ -calculus by the $\lambda\mu$ -calculus is smoother than the simulation of the $\lambda\mu$ -calculus by the $\bar{\lambda}\mu\tilde{\mu}$ -calculus. The first is obtained with the help of linear *reductions* whereas the second is obtained with the help of linear *expansions*.

This work can be extended in three ways. The first consists of proving the same results for the call-by-value evaluation of the $\lambda\mu$ -calculus defined in [Ong and Stewart, 1997]. The second consists of defining CPS translations to λ -calculus in order to complete [Curien and Herbelin, 2000]. The third consists of extending the type system to the other logical constants.

References

- Curien and Herbelin, 2000. Curien, P.-L. and Herbelin, H. (2000). The Duality of Computation. In *Proceedings of the International Conference on Functional Programming*.
- Dougherty et al., 2004. Dougherty, D. J., Ghilezan, S., and Lescanne, P. (2004). Characterizing strong normalization in a language with control operators. In *Proceedings of the 6th International ACM SIGPLAN Conference on Principles and Practice of Declarative Programming*, pages 155–166.
- Gentzen, 1934. Gentzen, G. (1934). Investigations into Logical Deduction. In Szabo, M., editor, *Collected Papers of Gerhard Gentzen*. North Holland.
- Griffin, 1990. Griffin, T. G. (1990). The Formulae-as-Types Notion of Control. In *Proceedings of the 17th ACM Symposium on Principles of Programming Languages*, pages 47–57. ACM Press.
- Ong and Stewart, 1997. Ong, L. and Stewart, C. (1997). A Curry-Howard foundation for functional computation with control. In *Proceedings of the 24th Annual ACM Symposium on Principles of Programming Languages*, pages 215–227. ACM Press.
- Parigot, 1992. Parigot, M. (1992). $\lambda\mu$ -calculus: an Algorithmic Interpretation of Classical Natural Deduction. In *Proceedings of International Conference on Logic Programming and Automated Deduction*, volume 624 of *Lectures Notes in Computer Science*, pages 190–201. Springer.